

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE Technical Papers		3. DATES COVERED (From - To)	
<div style="border: 1px solid black; border-radius: 50%; padding: 20px; text-align: center;"> <p>Please see attached</p> </div>				4. TITLE AND SUBTITLE	
				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
<div style="border: 1px solid black; border-radius: 50%; padding: 20px; text-align: center;"> <p>Please see attached</p> </div>				5d. PROJECT NUMBER 1011	
				5e. TASK NUMBER 0046	
				5f. WORK UNIT NUMBER 346204	
				8. PERFORMING ORGANIZATION REPORT	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				11. SPONSOR/MONITOR'S NUMBER(S) Please see attached	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
20030116 045					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT A	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Leilani Richardson
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code) (661) 275-5015

See'd. 20 Mar. 90
CJZ

MEMORANDUM FOR PR (In-House/Contractor Publication)
FROM: PROI (TI) (STINFO)

15 March 2000

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-TP-2000-049**
Larson, C. William; Harper, Jessica; Presilla-Marquez, J.D. (Schafer Corp.), "Matrix Isolation of Boron and Carbon Vapor. Control of Cluster Formation During Preparation and Annealing"

10th Internat'l IUPAC Conference on High Temperature Materials Chemistry (Statement A)
(Juelich, Germany, 10-14 April 2000) (Deadline: 31 March 2000)
and Seminars at Max Planck Inst., Univ of Dortmund, Univ. of Basel, Apr 4, 7, 18

**Matrix Isolation of Boron and Carbon Vapor.
Control of Cluster Formation During Preparation and Annealing.**

J. D. Presilla-Márquez, J. Harper, C. W. Larson
Propulsion Directorate
Air Force Research Laboratory
Edwards AFB, CA 93524-7680
Email: carl.larson@ple.af.mil

High Energy Density Matter (HEDM) Research Group
Pat Carrick (Chief), Jeff Sheehy (Group Leader), Greg Drake, Hi Young Yoo, Jeffrey Mills, Jerry Boatz, Jessica Harper,
Karl Christe, Mario Fajardo, Michael Tinnirello, Michelle DeRose, Paul Jones,
Txomin Presilla (Schafer Corporation) Peter Langhoff, Simon Tam, Suresh Suri, William Wilson,

**10th International IUPAC Conference on
High Temperature Materials Chemistry
Forschungszentrum Jülich
10-14 April 2000
Jülich, Germany**

**Matrix Isolation of Boron and Carbon Vapor.
Control of Cluster Formation During Preparation and Annealing.**

J. D. Presilla-Márquez, J. Harper, C. W. Larson
Propulsion Directorate
Air Force Research Laboratory
Edwards AFB, CA 93524-7680
Email: carl.larson@ple.af.mil

High Energy Density Matter (HEDM) Research Group
Pat Carrick (Chief), Jeff Sheehy (Group Leader), Greg Drake, Hi Young Yoo, Jeffrey Mills, Jerry Boatz, Jessica Harper,
Karl Christe, Mario Fajardo, Michael Tinnirello, Michelle DeRose, Paul Jones,
Txomin Presilla (Schafer Corporation) Peter Langhoff, Simon Tam, Suresh Suri, William Wilson,

**Universität Dortmund
Fachbereich Chemie
Anorganische Chemie
Dortmund, Germany
18 April 2000**

**Kinetics of formation of cyclic C_6 and cyclic C_8 and of
 $B_J C_{n-J}$ clusters ($J = 0, 1, 2$; $n = 3-11$) in solid argon**

J. D. Presilla-Márquez, J. Harper, C. W. Larson

**Propulsion Directorate
Air Force Research Laboratory
Edwards AFB, CA 93524-7680
Email: carl.larson@ple.af.mil**

High Energy Density Matter (HEDM) Research Group

Pat Carrick (Chief), Jeff Sheehy (Group Leader), Greg Drake, Hi Young Yoo, Jeffrey Mills, Jerry Boatz, Jessica Harper,
Karl Christe, Mario Fajardo, Michael Tinnirello, Michelle DeRose, Paul Jones,
Txomin Presilla (Schafer Corporation) Peter Langhoff, Simon Tam, Suresh Suri, William Wilson,

Institute für Physikalische Chemie

Universität Basel

Basel, Switzerland

7 April 2000

**Kinetics of formation of cyclic C_6 and cyclic C_8 and of
 $B_J C_{n-J}$ clusters ($J = 0, 1, 2$; $n = 3-11$) in solid argon**

J. D. Presilla-Márquez, J. Harper, C. W. Larson

**Propulsion Directorate
Air Force Research Laboratory
Edwards AFB, CA 93524-7680
Email: carl.larson@ple.af.mil**

High Energy Density Matter (HEDM) Research Group


Pat Carrick (Chief), Jeff Sheehy (Group Leader), Greg Drake, Hi Young Yoo, Jeffrey Mills, Jerry Boatz, Jessica Harper,
Karl Christe, Mario Fajardo, Michael Tinnirello, Michelle DeRose, Paul Jones,
Txomin Presilla (Schafer Corporation) Peter Langhoff, Simon Tam, Suresh Suri, William Wilson,

Max Planck Institute für Kernphysik

Heidelberg, Germany

5 April 2000

GOAL - 5% atoms in matrix



BCSyst4.xzg July 8, 1999 8:02:46 AM

Goal

Production of cryogenic HEDM with five mole percent atoms.

Objective

Characterization of source and quantitative analysis of $B_J C_{n-J}$

Approach

Production of HEDM by evaporation of boron with high-temperature graphite furnace and co-deposition of vapor with argon on a cold (10 K) surface


Identification and quantitative analysis of $B_J C_{n-J}$ species ($n \geq 3$, $J = 0$ to 2) by FTIR spectroscopy and *ab-initio* calculations.

Quantitative measurement of distributions of $B_J C_{n-J}$ species produced upon deposition and after annealing to a constant composition.

Determine absolute column densities (molecules cm^{-2}) from Beer's law:

$$\langle \rho_i \rangle = 2.303 A_{\text{exp}} / I_{\text{theory}}$$

Preparation




Annealing


<u>a0</u> 10 K	<u>a3</u> 32.5 K, 60 s	<u>a6</u> 40.0 K, 20 s
<u>a1</u> 27.5 K, 120 s	<u>a4</u> 35.0 K, 45 s	sublimation
<u>a2</u> 30.0 K, 90 s	<u>a5</u> 37.5 K, 20 s	rate ~ 1 $\mu\text{m}/\text{s}$

Precision matched pair of matrices


Green Matrix	$^{11}\text{B}/^{10}\text{B} = 80/20$	enhanced $^{11}\text{B}_j\text{C}_{n-j}$
Red Matrix	$^{11}\text{B}/^{10}\text{B} = 27/73$	enhanced $^{10}\text{B}_j\text{C}_{n-j}$



Survey spectra of precision matched matrices showing larger clusters B_JC_{n-J} , $n > 4$, $J = 0, 1, 2$ in original matrices and after three annealings.





Disappearance of BC_2 upon annealing





BC2rg3D Mar. 8, 2000 9:10:18 AM

Disappearance of BC_3 and C_4 upon annealing




BC3rg3D2.xvg Mar. 8, 2000 9:34:49 AM

Growth of B₂C₂ upon annealing




B2C2-3D2 Mar. 8, 2000 7:51:45 AM


Identification of 9 of the 16 isotopomers of linear BCCC in 5 matrices.



Four minimum energy geometries of B_2C_2 produce similar isotopomer fingerprints.
 Scale factor (S_F = measured frequency/theoretical frequency) of linear BCCB = 0.97250.



Identification of 7 isotopomers of the 10 isotopomers of BCCB in 5 matrices.



$B_J C_{n-J}$ Annealing Study

Peak by Peak

Detail


In first annealing C_4 and BC_3 remain constant and B_2C_2 grows from undetectable to $S/N \sim 100$. B_2C_2 seems inert.


All pentamers grow in first annealing. B_2C_3 grows most; C_5 and BC_4 evaporate most in 5th annealing.

Hexamers with boron grow most; evaporation of C_6 in 6th annealing is greatest.

In 5th annealing, $J = 0$ evaporates more than $J=1$, and $J = 1$ evaporates more than $J = 2$. A boron rich layer appears to be capped by a carbon rich layer.

In 6th annealing, where almost half the matrix sublimed, the $J = 2$ clusters evaporate the least. A boron rich layer appears to be capped by a carbon rich layer.





B32mol3.axg July 7, 1999 4:21:40 PM

Results and Discussion

Linear C₃, cyclic BC₂, and cyclic B₂C, constituted about 80% of the total observable boron and carbon in the initially deposited matrix, but B₃ was not observed. If B₃ were present, its concentration fell below the detection limit of the system. The measured trimer distribution in the initially formed matrices was $\rho(C_3) : \rho(BC_2) : \rho(B_2C) : \rho(B_3) \sim 1 : 1.5 : 0.5 : < 0.05$ (upper limit).

Statistical substitution of *J* boron atoms into an *n*-atom carbon cluster produces a distribution given by $\rho(B_J C_{n-J}) / \rho(C_n) = \{n(n-1)\dots(n-J+1)\} / J! [B/C]^J$. With the experimental B/C $\sim 1/3$, the statistical trimer distribution is

$$\rho(C_3) : \rho(BC_2) : \rho(B_2C) : \rho(B_3) \sim 1 : 1 : 0.33 : 0.03.$$

Agreement between distributions implies trimers form by random condensation of well-mixed atoms, uninfluenced by the relative energies of the trimers, the energies of their precursors, or preferential kinetics pathways that could otherwise distort the statistics.

Linear C₃ and cyclic BC₂, disappeared entirely when the matrices were repeatedly annealed to temperatures between 25 K and 35 K, but cyclic B₂C was inert.

Linear C₄ and BC₃ (BCCC) disappeared more slowly, and linear B₂C₂ (BCCB) grew to $\sim 95\%$ of its final value during the first annealing. Once formed, B₂C₂, like B₂C, was also inert to further reaction.

The sources of B_2C_2 are from condensation of atom plus trimer ($B + BC_2$ but not $C + B_2C$) or dimer + dimer ($BC + BC$ but not $B_2 + C_2$). Although BC was not observed, the upper limit of $\rho(BC)$ is larger than $\rho(B_2C_2)$ so that BC cannot be ruled out as a source of B_2C_2 .

The growth of B_2C_2 is conclusive evidence of the presence of BC and/or B in the originally deposited matrix in an amount at least as great as the growth of B_2C_2 .

Linear C_5 , BC_4 (BCCCC) and B_2C_3 (BCCCCB) and larger linear clusters (B_JC_{n-J} , $5 < n < 11$, $J = 0, 1, 2$), all grew upon annealing.

The sources of B_2C_3 are dimer + trimer ($BC + BC_2$ but not $B_2 + C_3$) and atom + tetramer ($B + BC_3$ but not $C + B_2C_2$).

Since $\rho(BC_2) \sim 5\rho(BC_3)$ in the initially deposited matrix, the $BC + BC_2$ source is dominant. Growth of B_2C_3 conclusively establishes the presence of BC in the matrix in an amount at least as great as the amount by which B_2C_3 grows.

Growth of BC_4 occurs primarily by $BC + C_3$ rather than $B + C_4$ or $C + BC_3$ because $\rho(C_3) \sim 10\rho(C_4)$ and $\rho(C_3) \sim 2\rho(BC_3)$. Growth of C_5 occurs by $C + C_4$ and $C_2 + C_3$, which establishes the presence of C and/or C_2 in the original matrix in an amount at least as great as C_5 growth.

Disappearance of triangular BC_2 requires breaking of one of its B-C bonds when one of its carbon atoms is attacked. The major reorganization of electronic energy involved in opening the ring appears to occur with little ($< \sim 3 \text{ kcal mol}^{-1}$) or no energy barrier, which makes this small molecule a candidate for an interesting *ab-initio* study of unusual reactivity at low temperature.

Conclusions

Annealing kinetics of disappearance of C_3 and BC_2 , and of appearance of B_2C , C_4 , BC_3 , B_2C_2 , C_5 , BC_4 , and B_2C_3 unequivocally establishes the presence of atoms and dimers in the originally deposited matrix.

$\sim 80\%$ or more of the initially deposited HEDM existed as atoms, dimers and trimers.

Molecules with two boron atoms are immune from radical attack and condensation during annealing.


Future Work

Continued development of source for production of higher flux beam of nearly pure boron atoms.


Map of "islands of stability" of pure boron HEDM; B_2 or B_3 may be the ultimate sink for atoms in the low temperature HEDM environment.

Determine reactivity of boron atoms with hydrogen during co-deposition.

Develop rapid condensation methodology to prevent reaction of B with H_2 .




Carbon Clusters




cgrasum2.axg July 8, 1999 7:13:47 AM

Carbon Clusters




cgrasumm.axg July 8, 1999 7:11:30 AM

Growth upon annealing - cC_6 and cC_8 - two fundamentals of C_{11}




MaCCn6.axg July 8, 1999 7:57:16 AM

Carbon Matrix (a) - Linear C_3 to C_{10} Clusters




MaCCn6.axg July 8, 1999 7:55:43 AM



Carbon Matrix FJ 32, JA03034


MaCCn5.axg July 8, 1999 7:56:39 AM

Theoretical Infrared Intensities of Linear C_n , DFT/B3LYP



TCNInte3.axg July 7, 1999 8:16:00 AM

Carbon cluster distributions



Most of the carbon condenses to cC₆ and cC₈ in all matrices

J218d8L.aug July 8, 1999 7:54:37 AM

Conclusions from Carbon HEDM Research

Quantitative analysis - Establishes HEDM density, distribution of carbon clusters, tracking of growth and decay of carbon clusters, carbon bookkeeping - quantification of "invisible carbon", C-atom and C₂.

Highest density matrix (equivalent C-atom density ~ 1 mole percent in argon) contained 40% "invisible" carbon (C, C₂), determined by tracking the growth of the "visible" (measurable) carbon to a constant composition after repeated annealing. Main product of condensation is cyclic C₆.

Yields of cyclic-C₆ are a factor of two larger than the combined yield of all other clusters in the fully condensed, highest density matrices. Cyclic-C₆ is the dominant condensation product.

Substrate must be shielded from oven to prevent condensation during deposition. Higher temperature oven places higher heat load on substrate, which promotes condensation.

Obtained higher density matrices by decreasing argon flux and maintaining oven flux. However, condensation was also increased.

Matrices produced with argon/5% H₂ caused nearly complete loss of C_{n+1} and C_{n+2} relative to C_{n+3}, suggesting that H₂ scavenges C-atoms efficiently during co-deposition.